Can integrative weaning index be a routine predictor for weaning success?

Sir,

We would like to bring you notice that the equation for integrative weaning index (IWI) has been mentioned incorrectly in the article by El-Baradey *et al.*;^[1] the correct equation is as follows:

IWI = $Crs \times SaO_2/(f/Vt)$.^[2,3]

Financial support and sponsorship Nil.

Conflicts of interest There are no conflicts of interest.

Sachin Gupta, Deepak Govil

Institute of Critical Care Medicine, Medanta - The Medicity, Gurgaon, Haryana, India

Correspondence: Dr. Sachin Gupta, Institute of Critical Care Medicine, Medanta - The Medicity, Gurgaon, Haryana, India. E-mail: dr_sachin78@yahoo.co.in

References

1. El-Baradey GF, El-Shmaa NS, Ganna SA. Can integrative weaning index be a routine predictor for weaning success? Indian J Crit Care Med 2015;19:703-7.

- Nemer SN, Barbas CS, Caldeira JB, Cárias TC, Santos RG, Almeida LC, et al. A new integrative weaning index of discontinuation from mechanical ventilation. Crit Care 2009;13:R152.
- Madani J, Saghafinia M, Nezhad H, Ebadi A, Ghochani A, Tavasoli F, et al. Validity of integrative weaning index of discontinuation from mechanical ventilation in Iranian ICUs. Thrita 2013;2:62-8.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.ijccm.org DOI: 10.4103/0972-5229.180056

predictor for weaning success?. Indian J Crit Care Med 2016;20:253-4.