

# Evaluation of the door-to-needle time for fibrinolytic administration for acute myocardial infarction

V. A. Masurkar, F. N. Kapadia, C. G. Shirwadkar, U. Shukla, P. Sood

## Abstract

**Background:** Fibrinolytic therapy has reduced mortality following acute myocardial infarction (AMI) with the major effect coming from early achievement of infarct-related artery patency. **Aim:** To evaluate the door-to-needle time for fibrinolytic administration for AMI and to identify factors associated with a prolonged door-to-needle time. **Materials and Methods:** Our study was a prospective audit of patients who were thrombolysed for AMI at our hospital from July 1, 2004 to March 15, 2005. All patients admitted with AMI, who were candidates for fibrinolysis, were included. We recorded the door-to-needle time. Whenever possible, we tried to find out the reason for prolonged door-to-needle time. **Results:** A door-to-needle time of <30 min could be achieved in 19 of our 35 patients (54.28%). Mean door-to-needle time was 45.25 min. **Discussion:** Although most guidelines recommend a door-to-needle time of less than 30 min, most hospitals fail to achieve this in most patients. A study conducted by Zed et al. at the Vancouver General Hospital showed that a door-to-needle time of less than 30 min was achieved in only 24.3%. The door-to-needle time achieved at our center was shorter. In most of our patients who were thrombolysed late, a delay in taking or interpreting an electrocardiogram was responsible. Transfer to the intensive care unit for thrombolysis also resulted in considerable delay. **Conclusions:** A door-to-needle time of less than 30 mins could be achieved in 19 of our 35 patients (54.28%). A significant number of AMI patients thrombolysed did not meet the guideline for door-to-needle time of less than 30 min.

**Key Words:** Acute myocardial infarction, Door-to-needle time, Fibrinolytic therapy

## Introduction

Fibrinolytic therapy has reduced mortality following acute myocardial infarction (AMI), with the major effect coming from early achievement of infarct-related artery patency. The Grampian region early anistreplase trial showed that delaying thrombolytic treatment by 1 h increases the hazard ratio of death by 20%, equivalent to the loss of 43/1000 lives within the next 5 years (95% CI

7–88,  $P = 0.012$ ). Delaying thrombolytic treatment by 30 min reduces the average expectation of life by approx 1 year.<sup>[1]</sup>

Thus, a short time to treatment interval must be considered as an adjunctive agent to fibrinolytic therapy. There are four components which determine the time between the onset of MI and achievement of reperfusion. (1) delay in seeking medical attention, (2) transport delays, (3) the door-to-needle time (the interval between the patient's arrival at the medical facility and the initiation of fibrinolytic therapy), and (4) fibrinolytic reperfusion time—the time between the administration of fibrinolytic therapy and the achievement of reperfusion.

### From:

Department of Critical Care Medicine, P.D. Hinduja National Hospital and Medical Research Centre, Mumbai

### Correspondence:

Dr. Vikram Masurkar, 4-6/ 19, Talmakiwadi, Javji Dadaji Marg, Tardeo, Mumbai – 400 007. E-mail: drmasurkar@hotmail.com

Free full text available from [www.ijccm.org](http://www.ijccm.org)

Efforts to reduce each of these components will lead to additive benefits in improving time to reperfusion and survival of patients with acute MI. The door-to-needle time is the easiest to modify.

### Materials and Methods

Our study was a prospective audit of patients who were thrombolized for AMI at our hospital from July 01, 2004 to March 15, 2005. We conducted a chart review of these patients. All patients admitted with AMI, who were candidates for fibrinolysis, were included. We recorded the following durations:

1. Time required to transfer the patient from the hospital entrance to the casualty.
2. Time taken to take an electrocardiogram (ECG) after arrival in the casualty.
3. Time needed to make a decision about thrombolysis after taking the ECG.

4. Duration between decision-making and actually starting thrombolysis.
5. Total door-to-needle time.

Whenever possible, we tried to find out the reason for prolonged door-to-needle time.

### Results

A door to needle time of less than 30 min could be achieved in 19 of our 35 patients. Table 1 shows the time taken to complete each step, which constitutes the door-to-needle time. Of the patients in whom there was a delay, in five patients the initial ECG showed subtle ST-segment changes which did not merit thrombolysis. Subsequent ECGs showed ST elevation. Two patients were transferred to the intensive care unit (ICU) for thrombolysis. The decision was taken by the casualty medical officer. One of our patients came to the hospital

**Table 1: Details of individual patients**

Door to casualty	Casualty to ECG	ECG to decision	Decision to thrombolysis	Door to needle	Reason/comment
3	2	<b>25<sup>a</sup></b>	5	<b>35</b>	Tr to ICU for thrombolysis
0	5	5	5	15	
0	5	5	0	10	
0	5	<b>30</b>	<b>15</b>	<b>50</b>	
2	3	5	5	15	
0	5	10	<b>20</b>	<b>35</b>	Tr to ICU for thrombolysis
0	5	5	<b>15</b>	25	
0	5	5	<b>10</b>	20	
0	5	5	5	15	
0	0	<i>80<sup>b</sup></i>	10	<i>90</i>	Subtle changes initially
2	0	5	5	12	
0	5	<i>35</i>	10	<i>50</i>	Subtle changes initially
2	0	0	10	12	ECG taken outside
0	5	10	5	20	
5	0	5	10	20	
0	5	0	5	10	
5	5	<b>35</b>	10	<b>55</b>	
0	10	10	6	26	
0	<b>40</b>	<b>20</b>	<b>15</b>	<b>150</b>	Delay in ECG, Tr to ICU
0	2	13	<b>15</b>	<b>30</b>	
2	<b>25</b>	10	10	47	Had VF. CPR for 5 min
2	5	0	<b>15</b>	22	
5	3	<b>25</b>	13	<b>33</b>	
2	0	0	10	<b>45</b>	ECG taken outside. ECG changes were not appreciated by CMO
0	10	<b>50</b>	10	<b>70</b>	
0	5	0	10	15	
0	5	<i>95</i>	5	<i>110</i>	Subtle changes initially
2	3	5	5	15	
2	3	<b>30</b>	5	<b>40</b>	
2	3	1	9	15	
0	5	<b>25</b>	5	<b>35</b>	
10	5	<i>285</i>	10	<i>310</i>	Subtle changes initially
2	10	<i>95</i>	15	<i>122</i>	Subtle changes initially
0	5	5	10	20	
5	0	5	5	15	
5	5	5	5	20	

<sup>a</sup>Entries in bold indicate unacceptable delays, <sup>b</sup>Entries in italic indicate delays with an explanation.

with an ECG taken by a general practitioner. The ECG showed an acute ST-elevation myocardial infarction (STEMI). However, the casualty medical officer did not identify these changes. One patient had a cardiac arrest owing to ventricular fibrillation (VF), which caused a delay. In seven patients, we could not identify any reason for a delay. Table 2 shows the door-to-needle time that was achieved in our study.

### Discussion

It has been proven that a shorter door-to-needle time results in better outcome. However, what is the standard of care? To address this question, Shuster and Dickinson<sup>[2]</sup> brought out recommendations for ensuring fibrinolytic therapy for AMI. They recommended early recognition of AMI symptoms by the public and health-care professionals, early access to emergency medical services, and early action by emergency-care providers in administering thrombolytic therapy (within 30 min after the patient's arrival at the emergency department). Grunfeld<sup>[3]</sup> responded to this article in the next issue of *Can Med Assoc J*. He mentioned that door-to-needle time of less than 30 min is probably an unrealistic goal. He also added that, to dogmatically adopt 30 min as the time interval during which all eligible patients are to receive thrombolytic therapy may well result in as many as half the patients receiving less than the recommended care.

In 2004, the AHA and ACC jointly brought out guidelines for the management of patients with STEMI. It was recommended that the delay from patient contact with the health-care system (arrival at the ED or contact with paramedics) to initiation of fibrinolytic therapy should be less than 30 min (level of evidence: B).<sup>[4]</sup>

The American College of Chest Physicians (ACCP) guidelines recommend that for patients with acute MI who are candidates for fibrinolytic therapy, the therapy should be administered within 30 min of arrival to the hospital or first contact with the health-care system (grade 1A).<sup>[5]</sup>

**Table 2: Door-to-needle time**

<i>n</i>	35
Mean door-to-needle time (min)	45.25
Door-to-needle time	
Less than 30 min	19/35 (54.28%)
30–40 min	4/35 (11.43%)
40–60 min	6/35 (17.14%)
>60 min	6/35 (17.14%)

Although most guidelines recommend a door-to-needle time of <30 min, most hospitals fail to achieve this in most patients. A study conducted by Zed *et al.*<sup>[6]</sup> at the Vancouver General Hospital showed that a door-to-needle time of <30 min was achieved in only 24.3%. Table 2 shows a comparison of our results with results from studies conducted at the Vancouver General Hospital and King Khalid University Hospital, Riyadh.<sup>[7]</sup>

Zed *et al.* found that shorter door-to-needle times were achieved when patients were thrombolized without a cardiology consult.<sup>[6]</sup> However, all patients at our center were thrombolized after a cardiology consult. Zed also noted that patients who arrived at the hospital during the night shifts were thrombolized faster. We found no such difference in our study.

In most of our patients who were thrombolized late, a delay in taking or interpreting an ECG was responsible. Transfer to ICU for thrombolysis also resulted in considerable delay. The above factors need to be looked into to improve door-to-needle time at our hospital.

### Conclusions

We could achieve a door-to-needle time of less than 30 min in 19 of our 35 patients (54.28%). A significant number of AMI patients thrombolized at our hospital do not meet the guideline for door-to-needle time of less than 30 min (Table 3). Factors associated with this should be addressed to improve the care of patients with AMI.

### References

1. Rawles JM. Quantification of the benefit of earlier thrombolytic therapy: Five-year results of the Grampian Region Early Anistreplase Trial (GREAT). *J Am Coll Cardiol* 1997;30:1181–6.
2. Shuster M, Dickinson G. Recommendations for ensuring early

**Table 3: Comparison of our results with results from other studies**

	Present study	Vancouver General Hospital	King Khalid University Hospital, Riyadh
<i>n</i>	35	140	271
Mean door-to-needle time (min)	45.25	58	95
Door-to-needle time			
Less than 30 min	19/35 (54.28%)	24.3%	
30–40 min	4/35 (11.43%)	24.3%	
40–60 min	6/35 (17.14%)	22.1%	
Greater than 60 min	6/35 (17.14%)	29.3%	

- thrombolytic therapy for acute myocardial infarction. *Can Med Assoc J* 1996;154:483–7.
3. Grunfeld AF. Reasonable “door-to-needle” time for thrombolytic therapy. *Can Med Assoc J* 1996;155:11–20.
  4. Antman EM, Anbe DT, Armstrong PW, Bates ER, Green LA, Hand M, *et al.* ACC/AHA guidelines for the management of patients with STEMI—executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2004;110:588–636.
  5. Menon V, Harrington RA, Hochman JS, Cannon CP, Goodman SD, Wilcox RG, *et al.* Thrombolysis and adjunctive therapy in acute myocardial infarction: the seventh ACCP conference on antithrombotic and thrombolytic therapy. *Chest* 2004;126:549S–75S.
  6. Zed PJ, Abu-Laban RB, Cadieu TM, Pursell RA, Filiatrault L. Fibrinolytic administration for acute myocardial infarction in a tertiary ED: Factors associated with an increased door-to-needle time. *Am J Emerg Med* 2004;22:192–6.
  7. Abba AA, Rahmatullah RA, Khalil MZ, Kumo AM, Wani BA, Ghonaim MA. Door to needle time in administering thrombolytic therapy for acute myocardial infarction. *Saudi Med J* 2003;24:361–4.

## Announcements

### 36th NATIONAL TRAUMA MANAGEMENT COURSE

We are glad to announce the 36th National Trauma Management Course-2006, which is scheduled to held at Auditorium, Sir Ganga Ram Hospital, Rajinder Nagar, New Delhi – 110060, on 16th - 17th February 2006, by the Dept. of Critical Care & Emergency Medicine, Sir Ganga Ram Hospital in association with Academy of Traumatology (India) and International Association for Surgery of Trauma & Surgical Intensive Care (IATSIC).

The objective of the course is to impart knowledge to delegates through teaching the techniques particularly applicable to trauma patients requiring immediate care. The course schedule comprises of lectures by renowned National & International faculties (South Africa, Finland, USA), demonstrations, skill stations (airway management, head injury etc.), case discussions & MCQ test. It is the only course of its kind available in India specifically designed taking into account Indian patients and the Indian scenario.

Since we are taking only 80 candidates for the course, please contact us for registration at the earliest.

To download the registration form and details of the course, visit the URL:

<http://sgrh.com/dept/criticare/critical.htm#Future%20Events>  
<http://sgrh.com/dept/criticare/36th%20NTMC-Delhi.pdf>

Thanking you,

**Dr. B.K. Rao**

Chairman – 36th National Trauma Management Course-2006  
Ex-President – ITACCS (Indian Chapter)