

# Indian Journal of CRITICAL CARE MEDICINE

Peer-reviewed, Official Publication of INDIAN SOCIETY OF CRITICAL CARE MEDICINE

**Indian Journal of Critical Care Medicine** is indexed/listed with Excerpta Medica/EMBASE, CAB Abstracts, Global Health, CINAHL Database, Cumulative Index to Nursing and Allied Health Literature, MANTIS, Health & Wellness Research Center, Health Reference Center Academic, InfoTrac One File, Expanded Academic ASAP, electronic databases of EBSCO, and Indian Science Abstracts.

Indian Journal of Critical Care Medicine is published quarterly (in March, June, September and December).

The information and opinions presented in Indian Journal of Critical Care Medicine reflect the views of the authors and not of the Society or the Editorial Board. Acceptance does not constitute endorsement by ISCCM.

#### All the rights are reserved.

Apart from any fair dealing for the purposes of research or private study, or criticism or review, no part of the publication can be reproduced, stored, or transmitted, in any form or by any means, without the prior permission.

Indian Journal of Critical Care Medicine and/or its publisher cannot be held responsible for errors or for any consequences arising from the use of the information contained in this journal. The appearance of advertising or product information in the various sections in the journal does not constitute an endorsement or approval by the journal and/or its publisher of the quality or value of the said product or of claims made for it by its manufacturer.

The Journal is printed on acid free paper.

#### EDITORIAL OFFICE

**Dr. Sandhya Talekar**  
Chief Editor, IJCCM  
Shree Medical Foundation, Prayag Hospital, 1247, Deccan Gymkhana, Pune 411004, India.  
Tel: 91 20 553 2812 / 553 2490  
Fax: 91 20 553 4448  
E-mail: editor@ijccm.org

#### PUBLISHED BY

**Medknow Publications**  
A-108/109, Kanara Business Center Off Link Rd, Ghatkopar (E), Mumbai - 400075, India.  
Phone: 91-22-6649 1818 / 1816,  
E-mail: publishing@medknow.com

#### WEBSITES

www.ijccm.org  
www.bioline.org.br/cm  
www.journalonweb.com/ijccm

#### CHIEF EDITOR

**Dr. Sandhya Talekar**  
sandhya\_talekar@yahoo.com

#### EXECUTIVE EDITOR

**Dr. Shirish Prayag**  
shirishprayag@gmail.com

#### ASSISTANT EDITORS

**Dr. R. K. Mani**  
rkmjs@vsnl.net

**Dr. Farhad Kapadia**  
fnkapadia@gmail.com

**Dr. J. V. Divatia**  
jdivatia@vsnl.com

#### MEMBERS OF EDITORIAL BOARD

**Dr. S. S. Iyer (Pune)**  
s\_ier@vsnl.com

**Dr. N. Ramakrishnan (Chennai)**  
icudoctor@gmail.com

**Dr. Atul Kulkarni (Mumbai)**  
kaivalyaak@yahoo.co.in

**Dr. Lalitha Pillai (Pune)**  
lalithapillai29@rediffmail.com

#### PEDIATRIC SECTION

**Dr. K. Chugh**  
gangaram@giasd01.vsnl.net.in

**Dr. Pravin Khilnani (New Delhi)**  
praveenk@nde.vsnl.net.in

**Dr. Sunit Singhi (Chandigarh)**  
drsinghi@glide.net.in

**Dr. Soonu Udani (Mumbai)**  
drudani@hotmail.com

#### NATIONAL ADVISORY BOARD

**Dr. M. B. Agarwal**

**Dr. H. S. Ballal**

**Dr. Dilip Karnad**

**Dr. Ram Rajagopalan**

**Dr. Manimala S. Rao**

**Dr. Subhash Todi**

**Dr. F. E. Udhwadia**

**Dr. R. S. Wadia**

#### INTERNATIONAL ADVISORY BOARD

**Dr. David Baker (France)**

**Dr. Laurent Brochard (France)**

**Dr. Geoffrey Dobb (Australia)**

**Dr. K. Guntupalli (USA)**

**Dr. Ken Hillman**

**Dr. Roop Kishen (UK)**

**Dr. Mitchell Levy (USA)**

**Dr. Ramesh Nagappan (Australia)**

**Dr. Michael Parr (Australia)**

**Dr. Gilbert Park (UK)**

**Dr. Ian Seppelt (Australia)**

**Dr. J. L. Vincent (Belgium)**

# Indian Journal of CRITICAL CARE MEDICINE

Official publication of Indian Society of Critical Care Medicine

Oct-Dec 2007

Volume 11, Issue 4

## C O N T E N T S

### Editorials

**It is not just the work - It is also the words**

– Murali Ramaswam \_\_\_\_\_ 169

**The ideal transfusion trigger in critically ill patients**

– M. B. Agarwal \_\_\_\_\_ 173

### Original Articles

**An increase in weight worsens the respiratory state and leads to intensive care unit re-admission**

– Yoshinori Matsuoka, Akinori Zaitso \_\_\_\_\_ 176

**A prospective cohort study on anemia and blood transfusion in critically ill patients**

– Naveen Manchal, S. Jayaram \_\_\_\_\_ 182

**Comparison of norepinephrine and dopamine in the management of septic shock using impedance cardiography**

– Sharad K. Mathur, Rajiv Dhunna, Arpan Chakraborty \_\_\_\_\_ 186

**Determination of intravascular volume status in critically ill patients using portable chest X-rays: Measurement of the vascular pedicle width**

– Nawal Salahuddin, M. Aslam, Ishtiaq Chishti, Shehla Siddiqui \_\_\_\_\_ 192

**Role of physiotherapists in intensive care units of India: A multicenter survey**

– Jithendra A. Kumar, Arun G. Maiya, Daphne Pereira \_\_\_\_\_ 198

### Review Article

**Postoperative pain and its management**

– Sona Chaturvedi, Amit Chaturvedi \_\_\_\_\_ 204

### Case Reports

**Rhabdomyolysis due to hair dye poisoning: An emerging threat**

– Krishnaswamy Sampathkumar, Yesudas Santhakumari Sooraj, Rajappannair Prabha Ajeshkumar, Amol Ramesh Mahaldar, Ramakrishnan Muthiah \_\_\_\_\_ 212

**Severe hyperkalemia with normal electrocardiogram**

– Sanjay Sharma, Harish Gupta, Meena Ghosh, Anantanarayan Padmanabhan \_\_\_\_\_ 215

**International Conference Calendar** \_\_\_\_\_ 218

**Author Index - 2007** \_\_\_\_\_ 000??

**Title Index - 2007** \_\_\_\_\_ 000??

# A prospective cohort study on anemia and blood transfusion in critically ill patients

Naveen Manchal, S. Jayaram

## Abstract

**Background:** The management of anemia and blood transfusion practices in the ICU have been a subject of controversy. **Aims:** The aims of this study were 1) To assess the prevalence of anemia and study the association of blood transfusion and mortality in critically ill patients. 2) To compare restrictive transfusion policy (Hb < 7 g/dl) and liberal transfusion policy. (Hb < 10 g/dl). **Settings and Design:** A matched cohort study was performed in a tertiary care teaching hospital. **Materials and Methods:** To study the association between blood transfusions and mortality, control patients were those who never received blood during ICU stay. They were selected according to the following matching criteria: Age ( $\pm 5$  years), sex, APACHE II score ( $\pm 5$  points), history of cardiac or renal disease and clinical diagnosis. **Statistical Analysis Used:** The Chi-square test. **Results:** The incidence of anemia is high in critically ill patients. Anemic patients had a longer duration of stay in the ICU. There is an association between blood transfusion and higher mortality in critically ill patients. A restrictive transfusion policy was associated with lesser mortality. **Conclusions:** Anemia is associated with increased morbidity reflected by the increased duration of stay in the ICU. Blood transfusion is associated with increased mortality and a restrictive transfusion policy is associated with increased survival.

**Key words:** Anemia, blood transfusion, critically ill, mortality

## Introduction

As the entire practice of medicine has evolved in the past few decades, so have transfusion practices. Two decades ago, a hematocrit between 0.20 and 0.25 was considered an urgent indication for transfusion, but at the turn of this century, maintaining a hematocrit at this level is considered to be “best-practice medicine”.<sup>[1]</sup>

Nearly two thirds of patients admitted to an ICU have hemoglobin levels of 10 g/dl or lower.<sup>[1]</sup> Although intraoperative blood loss and gastro intestinal hemorrhage contribute to these statistics and are a frequent reason

for administering transfusions, only 40% of transfusions administered were because of acute blood loss.

Although anemia is apparently tolerated in most patients, particularly those who are young /relatively healthy, the ICU population must be thought of differently. Anemia in the ICU may be due to acute blood loss, phlebotomy or to the presence of inflammatory disease. The risks of blood transfusions are many. Nonetheless, hemoglobin levels at or above 10 g/dl may be important for oxygen delivery to vital organs, especially in critically ill patients with increased oxygen demands.<sup>[1]</sup> The appropriate transfusion trigger for critically ill patients in this setting remains unknown.<sup>[6,8,16-17]</sup>

This study aims to assess the prevalence of anemia in critically ill patients and studies the association of blood transfusion and mortality in critically ill patients. It also

**From:**  
Department of Medicine, J. J. Group of Hospitals, Mumbai - 400 008, India

**Correspondence:**  
Dr. Naveen Manchal, No. 278, 5<sup>th</sup> Main, 5<sup>th</sup> Cross, Micokgnss Layout, Arekere, Bangalore - 76, Karnataka, India. E-mail: drsoda79@yahoo.com

Free full text available from [www.ijccm.org](http://www.ijccm.org)

compares restrictive transfusion policy (Hb < 7 g/dl) and liberal transfusion policy (Hb < 10 g/dl).

### Materials and Methods

A total of 100 patients were included in the study. A matched cohort study was performed. The complete blood count was done every week and patients were followed for 21 days or till death/ transfer from the ICU. The following standards were used for case identification:- Hb less than 12 g/dl (males). Hb less than 10 g/dl (females) [any single episode during ICU stay].

#### Inclusion criteria

All patients admitted in the intensive care unit during the study period in the age group of 15-65 years.

#### Exclusion criteria

1. Patients with a known hematological disorder
2. Hematological malignancies
3. Patients on chemotherapy
4. HIV positive patients

#### Matching and selection of control patients

To study the association between blood transfusions and mortality, control patients were those who never received blood during ICU stay. They were selected according to the following matching criteria:

- Age (±5 years)
- Sex
- APACHE II score on first day of ICU admission (± 5 points)
- History of cardiac disease (ischemic heart disease, congestive heart failure)
- History of renal disease (acute or chronic renal failure)
- Clinical diagnosis
- The Chi-square test was used for statistical significance and  $P < 0.05$  was considered significant at 95% confidence interval.


### Results

The study cohort of 100 patients had an average APACHE score of 16.2. The overall mortality was 46% and the age-wise distribution of mortality was as shown in Table 1.

1. Out of 100 patients 68 were found to be anemic during the study period. The mean Hb level was 9.76 g/dl. The predominant peripheral smear picture


**Table 1: Age wise distribution of mortality**

Age (years)	Number of deaths
15-24	6
25-34	8
35-44	11
45-54	12
55-64	9



Bar diagram showing age-wise distribution of mortality

**Table 2: The predominant peripheral smear picture was microcytic hypochromic**



was of microcytic hypochromic type. The others were normocytic normochromic with 3% showing a leukemoid reaction [Table 2].

2. Sepsis<sup>[4]</sup> was the commonest clinical setting associated with anemia. The others being gastrointestinal bleed, chronic renal failure and postoperative blood loss [Table 3].
3. Out of the 100 patients studied, 38 patients were given blood transfusion and matching control patients were identified for 36 patients. The mortality in the

transfusion group was 58.33% when compared to 25.0% in the non transfusion group ( $P$  0.005). The attributable mortality was 33.33% with a relative risk of 2.33 [Table 4].

4. The mortality rate in those transfused at  $Hb < 7$  g/l was 10% when compared to 70% mortality rate in those transfused at  $Hb < 10$  g/l ( $P$  0.005) [Table 5]
5. The duration of stay in the ICU was doubled in anemic patients [Table 6].

### Discussion


The study shows that nearly two-thirds of patients admitted to an ICU have Hb levels less than 10 g/l. Anemia may be caused by decreased production of RBCs, nutritional deficiencies, inadequate endogenous erythropoietin production, renal failure and diagnostic blood sampling.<sup>[7,14,15,18,22]</sup>

**Table 3: Causes of anemia identified in the study**

Aetiology	No. of patients
GI. Bleed	8
CRF	9
Sepsis	33
Blood loss due to internal hemorrhage, post operative etc.	9
Protozoal infections (malaria, leptospirosis)	8

**Table 4: Association between blood transfusion and mortality**

	No. of deaths	No. of survivors
Total number transfused	21	15
Number of patients not transfused	9	27



**Pie diagram showing mortality in transfused and non transfused groups**

**Table 5: The results comparing two strategies of blood transfusion are as follows: out of the 21 patients transfused, matching controls were found for 10**

	Restrictive transfusion (Trigger $< 7$ g/L)	Liberal transfusion (Trigger $< 10$ g/L)
Mortality	10.0%	70.0% ( $P$ 0.005)

**Table 6: Duration of ICU stay**


	Anemia	Normal Hb
Average duration of stay in ICU (days)	8.72	4.10

As anemia progresses in otherwise healthy persons, compensatory mechanisms are recruited and a certain degree of anemia may be tolerated. These mechanisms may not operate efficiently or at all in critically ill patients. Under various preconditions anemia may be associated with increased mortality of the critically ill. The same is true with blood transfusion.<sup>[13,19]</sup>

The complications of blood transfusions include volume overload, febrile reactions and fatal hemolytic reactions which may all contribute to increased mortality.<sup>[5,9-11,20,21]</sup> The anemia and blood transfusions study included 3534 patients from 146 Western European ICUs. Both ICU and overall mortality rates were significantly higher in patients who had vs who had not received a transfusion (ICU rates: 18.5% vs 10.1% respectively;  $P < 0.001$ ).<sup>[2]</sup>

The concept of optimal Hb concentration has been challenged. Wilkerson and colleagues using a paralyzed, anesthetized, normovolemic anemic primate model, noted that compensatory mechanism for low Hb concentration did not occur until the Hb concentration fell below 7 g/dl.<sup>[14]</sup>

The study by Hebert *et al.*,<sup>[12]</sup> showed that a restrictive transfusion policy ( $Hb < 7$  g/dl) had better overall outcomes than when the transfusion trigger was more liberal ( $Hb < 10$  g/dl).<sup>[3]</sup> The decision to transfuse or not must be taken after weighing up the potential consequences of anemia against blood transfusion associated risks. Every single patient's ability to compensate for anemia is different. The question of whether a potential transfusion is capable of affecting the clinical outcome of a given patient is related to the question whether tissue hypoxia is in fact prevailing at a certain Hb concentration.



**Clinical settings associated with anemia**

## Acknowledgement

Dr. Aditi Manchal for her valuable comments and help with the manuscript.

## References

1. Pearl RG, Pohlman A. Understanding and managing anemia in critically ill patients. *Crit Care Nurse* Dec 2002;Suppl:1-14.
2. Vincent JL, Baron JF, Reinhart K, Gattinoni L, Thijs L, Webb A, *et al.* Anemia and blood transfusion in critically ill patients. *JAMA* 2002;288:1499-507.
3. Hebert PC, Well G, Martin C, Tweeddale M, Marshall J, Blajchman M, *et al.* Variation in red cell transfusion practice in the intensive care unit: A multicentre cohort study. *Crit Care* 1999;3:57-63.
4. American College of Chest physicians/Society of Critical Care Medicine Consensus Committee: Definitions for sepsis and organ failures and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992;20:864-74.
5. Carson JL, Duff A, Poses RM, Berlin JA, Spence RK, Trout R, *et al.* effect of anemia and cardiovascular disease on surgical mortality and morbidity. *Lancet* 1996;348:1055-60.
6. Carson JL, Hill S, Carless P, Hebert P, Henry D. Transfusion triggers: A systematic review of the literature. *Transfus Med Rev* 2002;16:187-99.
7. Warrel DA, Lox TM, Firth JD, Benz EJ Jr. *Oxford text of medicine* 4<sup>th</sup> ed. 2003. p. 1219-27.
8. Drews RE. Critical issues in hematology: Anemia, thrombocytopenia, coagulopathy, and blood product transfusions in critically ill patients. *Clin Chest Med* 2003;24:607-22.
9. Edna TH, Bjerkeset T. Association between blood transfusion and infection in injured patients. *J Trauma* 1992;33:659-61.
10. Stephan F, Montblanc Jd, Cheffi A, Bonnet F. Thrombocytopenia in critically ill surgical patients: A case-control study evaluating attributable mortality and transfusion requirements. *Crit Care* 1999;3:151-8.
11. Goodnough LT, Brecher ME, Kanter MH, AuBuchon JP. Transfusion medicine: First of two parts-blood transfusion. *N Engl J Med* 1999;340:438-47.
12. Herbert PC, Well G, Tweeddale M, Martin C, Marshall J, Pham B, *et al.* Does transfusion practice affect mortality in critically ill patients? *Am J Respir Crit Care Med* 1997;155:1618-23.
13. Jesse B Hay, Gregory A Schmidt, Laurence DH Wood. *Principles of critical care* 1992. p. 1807-17.
14. Civetta JM, Taylor RW, Kirby RR. Transfusion therapy. *Crit Care* 1996;vol 3<sup>rd</sup> Edition:639-52.
15. Kasper, Braunwald, Fauci, Hauser, Longo, Jameson. *Transfusion biology and therapy. Harrison's principles of internal medicine.* 16<sup>th</sup> ed. 2005. p. 662-7.
16. Practice guidelines for blood component therapy: A report by the American Society of Anesthesiologists task force on blood component therapy. *Anesthesiology* 1996;84:732-47.
17. Schroder ML. Principles and practice of transfusion medicine. *In: Wintrobe's Clinical Hematology.* Lee GR, Foerster J, Lukens J, *et al*, editors. Williams and Wilkin: Baltimore; 1998. p. 817-74.
18. Shoemaker, Ayers, Grenvik, Holbrook, Thompson: *Textbook of critical care.* p. 903-18.
19. Evans TW, (2001) Hemodynamic and metabolic therapy in critically ill patients, *NEJM* 345:1368-77.
20. Thurer RL. Blood transfusion in cardiac surgery. *Can J Anesth* 2001;48:S6-12.
21. Vamvakas ES, Moore SB, Cabanela M. Blood Transfusion and septic complications after hip replacement surgery. *Transfusion* 1995;35:150-6.
22. Wallace EL, Churchill WH, Surgenor DM, Cho GS, McGurk S. Collection and transfusion of blood components in the United States, 1994. *Transfusion* 1998;38:625-36.

**Source of Support:** Nil, **Conflict of Interest:** None declared.