

# Clinical Profile and Outcome of Dengue Fever in Multidisciplinary Intensive Care Unit of a Tertiary Level Hospital in India

Mahesha Padyana<sup>1</sup>, Sunil Karanth<sup>2</sup>, Shriram Vaidya<sup>3</sup>, Justin Aryabhat Gopaldas<sup>4</sup>

## ABSTRACT

**Background:** India is one of the seven identified countries in South-East Asia region regularly reporting dengue fever (DF)/dengue hemorrhagic fever (DHF) outbreaks. Even though the dengue prodrome and evolution of illness are most often similar in many patients, progress and outcome may differ significantly depending on the severity of illness as well as treatment instituted. We studied the clinical manifestations, outcome and factors predicting mortality of serology confirmed dengue fever cases admitted in Multidisciplinary Intensive Care Unit (MICU) of a high acuity healthcare facility in India.

**Methodology:** All patients with serology proven dengue fever admitted to MICU between 1st July 2015 and 1st December 2015 were included in the study. Clinical presentation, laboratory findings, severity of illness scores and outcome were recorded.

**Results:** Majority of the patients (58.4%) belonged to 21–40 year age group. Hepatic (96.8%) followed by hematological (79.2%) involvement were the most common findings. CNS involvement observed among 27%. Survival to hospital discharge was 78.9%. Respiratory and gastrointestinal system involvement was associated with increased mortality. Acute respiratory distress syndrome (ARDS), acute kidney injury (AKI) and shock were the clinical syndromes associated with mortality. Serum lactate, aspartate transaminase (AST) and alanine transaminase (ALT) were significantly elevated among non survivors. Significant difference in sequential organ failure assessment (SOFA) and acute physiology and chronic health evaluation (APACHE) scores was also observed among survivors and non survivors.

**Conclusion:** Organ system involvement and higher disease severity scores are strong predictors of mortality. High index of suspicion for atypical manifestations of dengue is warranted.

**Keywords:** APACHE, Dengue fever, SOFA, Severe Dengue, Transaminitis

*Indian Journal of Critical Care Medicine* (2019): 10.5005/jp-journals-10071-23178

## INTRODUCTION

Dengue is a vector born tropical viral fever spread by *Aedes aegypti* and *Aedes albopictus* mosquitoes. Incidence of dengue is rising in India from 6.34 per million population between 1998 and 2009 to 34.81 per million population between 2010 and 2014.<sup>1</sup> Presentation of dengue fever can vary from malaise, fatigability as a part viral prodrome to shock and multi organ dysfunction syndrome as a part of severe illness.<sup>2</sup> Outcomes of patients with dengue fever admitted to critical care units have been studied less often. Prognostic factors which determine the clinical outcome of critically ill patients with dengue fever remain unclear.<sup>3</sup>

## METHODOLOGY

Retrospective single center study which included patients admitted to MICU with serology proven<sup>4</sup> dengue fever with warning signs and severe dengue between 1st July and 1st December 2015 NS1-Antigen Strip<sup>®</sup> ELISA (BioRad), IgM and IgG- Capture ELISA Panbio<sup>®</sup> (Alere<sup>TM</sup>) were used for dengue serology testing. Patient management was as per WHO, Surviving Sepsis Campaign guidelines and established standards in intensive care practice. Demographics including age and gender distribution, signs and symptoms, SOFA and APACHE II scores were recorded. Outcome details including organ involvement and survival to hospital discharge were recorded. Statistical Analysis done using SPSS version 13.0 used for data analysis. Data analyzed by frequency,

<sup>1–4</sup>Department of Critical Care Medicine, Manipal Hospital, Bangalore, Karnataka, India

**Corresponding Author:** Mahesha Padyana, Department of Critical Care Medicine, Manipal Hospital, Bangalore, Karnataka, India, Phone: 9448843933, e-mail: padyana@gmail.com

**How to cite this article:** Padyana M, Karanth S, Vaidya S, Gopaldas JA. Clinical Profile and Outcome of Dengue Fever in Multidisciplinary Intensive Care Unit of a Tertiary Level Hospital in India. *Indian J Crit Care Med* 2019;23(6):270–273.

**Source of support:** Nil

**Conflict of interest:** None

percentage, mean ( $\pm$ SD)/ Median (Inter Quartile Range- IQR). Logistic regression analysis for determination of factors predicting mortality.

## RESULTS

Total admissions to MICU during study period was 1,170 out of which patients with dengue fever were 96. Median length of ICU stay among severe dengue was 10 (IQR 14-8), dengue with warning sign was 4 (IQR 5-3) and dengue with no warning signs was 3 (IQR 5.75-2). Out of 96 patients with dengue, 47 (48.9%) were males and 49 were (51.1%) females. Patients belonged to age group between 21 years and 40 years were 58.4% (Table 1). Fifty-one patients had associated conditions like diabetes mellitus (18.5%), hypertension

**Table 1: Demographics**

		Number	%
Age (in years)	<20	8	8.3%
	21 - 40	56	58.4%
	41 - 60	14	14.5%
	>60	18	18.8%
Sex	Males	47	49.0%
	Females	49	51.0%

**Table 2: Clinical features**

Symptoms and signs	Frequency	%
Fever	91	94.8%
Headache	39	40.6%
Bodyache	50	52.1%
Generalised weakness	75	78.9%
Loss of appetite	62	64.6%
Vomiting	56	58.3%
Abdominal pain	40	41.7%
Loose stools	20	20.8%
Breathlessness	55	57.3%
Bleeding manifestation	30	31.3%
Petichiae/echymosis	38	39.6%
Jaundice	16	16.7%
Altered sensorium	26	27.1%

(14.4%), pregnancy (9.3%). Predominant symptom was fever (94.8%) followed by warning signs like vomiting (58.3%), breathlessness (57.3%), abdominal pain (41.7%), bleeding manifestations (31.3%) and altered sensorium (27.1%) (Table 2).

Predominant organ involvement was hepatic (transaminitis, jaundice) followed by gastrointestinal (abdominal pain, distension, loose stools, vomiting, ascitis) systems. Forty-four percent patients had gall bladder wall edema on ultrasound evaluation. Hematological involvement was predominantly thrombocytopenia and raised hematocrit. Respiratory system involvement had ARDS and Pleural effusion. CNS manifestations included intracranial bleed, cerebral infarction and cerebellitis (in one patient). Patients with refractory shock requiring vasopressor initiation were 30.2% (Fig. 1). An average of 2 units of single donor platelet transfusion requirement per patient was noted in our study (Table 3).

Overall mortality was 21.1%. Among non survivors 12 (60%) females of which 8 belonged to 21-40 year age group. Median APACHE II score at 24 hours among survivors was 6.0 and non-survivors 17.5 ( $p < 0.01$ ). SOFA score at 48 hours among survivors and non survivors was 6 and 20, respectively ( $p < 0.01$ ) (Tables 4, 5 and 6).

Total of 42 (43.8%) patients needed mechanical ventilation. Twenty-two of 76 patients (28.9%) who survived required mechanical ventilation. Acute kidney injury was noted among 34 (35.4%) patients of which 24 (25%) needed either CRRT or SLEDD. Of all, 18.4% of survivors required renal replacement therapy and 29 patients (30.2%) required vasopressor support during ICU stay.


Secondary sepsis was noted in 16.5% of patients. Half of these were blood stream infections. Predominant organisms responsible are gram-negative (*Klebsiella pneumoniae* and *Acinetobacter baumannii*). Ten (13.1%) patients required tracheostomy for liberation of mechanical ventilation due to ICU acquired weakness (5.2%) and persistent neurological abnormalities like hemiparesis, ataxia, dysarthria etc.

**Table 3: Transfusion pattern**

Blood product	Mean no of units/patient
Single donor platelet	2
Packed red blood cell	0.7
Fresh frozen plasma	1.2

**Table 4: Mortality**

	Frequency	Percentage (%)
Survivors	76	78.9
Non survivors	20	21.1



**Fig. 1: Organ involvement in patients during dengue fever**

## DISCUSSION

Our study looked into epidemiological factors, organ involvement and mortality during ICU stay.

In our study, we noted that 50 patients (52.1%) had dengue with warning signs and 41 patients (42.7%) had severe dengue and only 5 patients (5.2%) had dengue without warning signs based on WHO classification.<sup>4</sup> We observed that the mean age of our cohort of patients was 37 years. ICU requirement was significantly high among age group of 21-40 (58.4%). Similarly, in a study by Juneja et al. 39.6 year was the mean age of patients requiring ICU admission with dengue fever suggesting that younger age group is at higher risk compared to older.<sup>5</sup> Vomiting, abdominal pain and bleeding manifestations were common warning signs which warranted ICU transfer, similar to observations in previous studies.<sup>3,6</sup>

Multiorgan involvement in dengue is well known. In our study we observed Hepatic involvement among 96.8%. AST and ALT were significantly high among non survivors. Similarly in a study by Nayak et al liver involvement was noted among 97.33%, AST and ALT were significantly high among dengue hemorrhagic fever and dengue shock syndrome group.<sup>7</sup>

Respiratory and gastrointestinal involvements had increased risk of mortality. AKI and moderate to severe ARDS (with PF ratio  $< 200$ ) was significantly high among non survivors. Similarly Shock and Lactates were also noted to be significantly high among non survivors. Study by Thanachartwet et al showed significantly high lactates among patients with organ failure and shock.<sup>8</sup> In a study by Chen et al moderate ARDS (PF Ratio  $< 200$ ) followed by shock (SBP  $< 90$  mm Hg or MAP  $< 65$ ) and multiorgan failure was significant

**Table 5:** Factors associated with mortality

Variable	Survivors (n = 76)	Non survivors (n = 20)	OR (IQR)	p value
Jaundice	8 (10.5%)	8 (40%)	5.66 (1.78-18.0)	0.003
ARDS(PaO <sub>2</sub> /FiO <sub>2</sub> <200)	22 (28.9%)	17 (85%)	13.9 (3.7-52.3)	<0.01
Acute kidney injury (AKI)	15 (19.7%)	19 (95%)	77.26 (9.56-624)	<0.01
Shock	16 (21%)	19 (95%)	71.3 (8.86-574)	<0.01
Central nervous system	20 (26.3%)	8 (40%)	1.86 (0.66-5.23)	0.235
Respiratory	48 (63.2%)	19 (95%)	11.08 (1.4-87.3)	<0.05
Gastrointestinal	45 (59.2%)	19 (95%)	13.1 (1.67-103)	<0.05

**Table 6:** Evaluation of variables in survivors and non survivors

Variable	Survivors (n = 76)	Non survivors (n = 20)	p value
Lactate (IQR)	3.0 (20-4.8)	13.4 (9.45-7.5)	<0.01
Aspartate Transaminase (IQR)	299.5 (148-1098.5)	4936 (2431.5-14340.3)	<0.01
Alanine Transaminase (IQR)	163 (83.75-469.5)	1412 (613-3832.75)	<0.01
Base Excess	-3.75 (-1.42-6.57)	-21.9 (-18.2- -24.0)	<0.01
Platelets (mean ±SD)	22493.4 (24405.3)	19200 (15949.3)	0.864
Total count (mean ±SD)	8258 (6032)	(13453 (10799.9)	0.095
Hematocrit (mean ±SD)	41.5 (9.84)	43.2 (9.54)	0.332
Serum albumin (g/dL), (mean ±SD)	2.67 (0.62)	2.44 (0.7)	0.264
APACHE median (IQR)	6.0 (4.0-9.0)	17.5 (9.5-35.75)	<0.01
SOFA 48 hours median (IQR)	6 (3.0-20.0)	20 (18-22)	<0.01

among non survivors.<sup>3</sup> A study by Jain et al. had described liver and renal involvement, ARDS, Shock requiring vasopressor, metabolic acidosis as predictors of mortality.<sup>9</sup> A study done in Cuttack, India had showed predominant organ failure was coagulation (48%). Respiratory failure was 19% and cardiovascular failure was 8%.<sup>10</sup>

Neurological involvement may occur without associated bleeding manifestations, rash or thrombocytopenia. Neurological complications are reported to be rare in dengue as it is a non-neurotropic virus.<sup>11</sup> However we observed that many patients (27.1%) were presented with altered sensorium as initial manifestation either due to primary CNS involvement (encephalopathy, encephalitis or intracranial bleed) or secondary to profound shock. Karoli et al described neurological involvement upto 14% in patients with dengue fever in the form of encephalopathy, hypokalemic periodic paralysis, myositis and Guillain-Barre syndrome. Another study by Nimmagadda et al placed incidence of neurological manifestations at 2% encephalitis or meningitis.<sup>12,13</sup> In our study, three patients had infarct, two sustained intracranial hemorrhage, one had Posterior Reversible Encephalopathy Syndrome (PRES) and other was found to have cerebral abscess with hydrocephalus (overall incidence 8.3%). One of the patients with multiple infarcts had cerebrospinal fluid polymerase chain reaction (PCR) positive for dengue. Recently published data from Taiwan demonstrated that altered sensorium is an independent risk factor for mortality.<sup>14</sup> However in our study, we did not find any significant association between altered sensorium and mortality.

Mortality has been shown to be high among age group of >60 years in prior studies.<sup>3</sup> Our study had younger patients between 21 years and 40 years requiring ICU and mortality was noted to be high among females between the age group of 21–40 years. In a retrospective cohort study from Brazil mortality of severe dengue was noted to be 3.1% over a period of 14 years. Age >55 year was associated with increased mortality.<sup>15</sup> In a Chinese study, female sex and age group 30–39 years was associated with increased risk

of dengue hemorrhagic fever and mortality especially with dengue 2 serotype<sup>6</sup> Pang et al noted that adult patients belonging to 50 to 59 year age group or patients with diabetes had 5 times increased risk for ICU admission and increased mortality compared to patients who are younger than 30 year and without diabetes.<sup>16</sup> Another study done in Singapore between 2007 and 2008 during dengue sero-type 2 epidemic observed that female sex, age group 30–49 years and presence of diabetes or hypertension were associated with increased risk of dengue hemorrhagic fever.<sup>17</sup> However in our study, diabetes was not associated with significant increase in mortality.

There are certain limitations in our study. Interpretation of this study should be done bearing in mind that it is a retrospective single center study involving patients admitted in the intensive care unit. However, this study sheds light onto the factors which could predict mortality among patients admitted with dengue to critical care unit.

## CONCLUSION

Lactatemia, transaminitis, higher disease severity scores (APACHE II and SOFA), Shock, AKI and PaO<sub>2</sub>/FiO<sub>2</sub> ratio less than 200 were associated with increased mortality. We advise high index of suspicion for atypical CNS manifestations in patients admitted with dengue fever to intensive care unit.

## REFERENCES

- Laul A, Laul P, Merugumala V, Pathak R, Miglani U, Saxena P. Clinical Profiles of Dengue Infection during an Outbreak in Northern India. *J Tropical Med*. 2016; 1-7. <http://dx.doi.org/10.1155/2016/5917934>
- Mutheni SR, Morse AP, Caminade C, Upadhyayula SM. Dengue burden in India: recent trends and importance of climatic parameters. *Emerg Microbes Infect*. 2017; 6(8): e70. doi: 10.1038/emi.2017.57
- Chen CM, Chan KS, Yu WL, Cheng KC, Chao HC, Yeh CY et al. The outcomes of patients with severe dengue admitted to intensive care units. *Medicine* 2016;95:31. doi: 10.1097/MD.00000000000004376

4. World Health Organization (WHO): Dengue: Guidelines for Diagnosis, Treatment, Prevention and Control. Geneva: WHO; 2009.
5. Juneja D, Nasa P, Singh O, Javeri Y, Uniyal B, Dang R: Clinical profile, intensive care unit course, and outcome of patients admitted in intensive care unit with dengue. *J Crit Care* 2011; 26(5):449–452. doi: 10.1016/j.jcrc.2011.05.007
6. Ong A, Sandar M, Chen MI, Sin LY. Fatal dengue hemorrhagic fever in adults during a dengue epidemic in Singapore. *Int J Infect Dis* 2007;11: 263-267. DOI: 10.1016/j.ijid.2006.02.012
7. Nayak J, Behera S, Swain SK, Panda SR. A study of multiorgan dysfunction in patients with dengue and its clinicohematological correlation with severity. *Asian J Pharm Clin Res*. 2017;10(2):218–221.
8. Thanachartwet V, Desakorn V, Sahassananda D, Jittmittraphap A, Areemitr NO, Osothsomboon S et al. Serum Procalcitonin and peripheral venous lactate for predicting Dengue Shock and/or organ failure: A prospective observational study. *PlosNegl Trop Dis*. 2016; 10(8): e 0004961. <https://doi.org/10.1371/journal.pntd.0004961>
9. Jain S, Mittal A, Sharma SK, Upadhyay AD, Pandey RM, Sinha S et al. Predictors of Dengue-Related Mortality and Disease Severity in a Tertiary Care Center in North India. *Open Forum Infectious Diseases*. 2017; 4 (2), 1-8. doi: 10.1093/ofid/ofx056.
10. Routray SS, Mohanty DP, Mishra D. Clinical profile, Intensive care Unit course and outcome of patients admitted in intensive care unit with dengue. *Indian J Clin Anaesth*. 2016; 3(1): 33-35
11. Madi D, Achappa B, Ramapuram JT, Chowta N, Laxman M, Mahalingam S. Dengue encephalitis- A rare manifestation of dengue fever. *Asian Pac J Trop Biomed* 2014; 4(suppl 1):S 70- S72. doi: 10.12980/APJTB.4.2014C1006.
12. Karoli R, Fatima J, Siddiqi Z, Kazmi KI, Sultania AR . Clinical profile of dengue infection at a teaching hospital in North India. 2010; 2010–2013.
13. Nimmagadda SS , Mahabala C , Boloora A , Raghuram PM , Nayak AU. Atypical Manifestations of Dengue Fever (DF) – Where Do We Stand Today? *J Clin Diag Res*. 2014; 8(1): 71-73. doi: 10.7860/JCDR/2014/6885.3960
14. Yeh CY, Chen PL, Chuang KT, Shu, Chien YW, Perng GC et al. Symptoms associated with adverse dengue fever prognoses at the time of reporting in the 2015 dengue outbreak in Taiwan. *PlosNegl Trop Dis*. 2017; 11(12): e 0006091. doi: 10.1371/journal.pntd.0006091
15. Pinto RC, Castro DBd, Albuquerque BCd, SampaioVdS, PassosRAd, Costa CFd, et al. Mortality Predictors in Patients with Severe Dengue in the State of Amazonas, Brazil. *PLoS ONE*. 2016;11(8): e0161884. <https://doi.org/10.1371/journal.pone.0161884>
16. Pang J, Thein TL, Leo YS, Lye DC. Early clinical and laboratory risk factors of intensive care unit requirement during 2004–2008 dengue epidemics in Singapore: a matched case–control study. *BMC Infect Dis* 2014; 14:649. doi: 10.1186/s12879-014-0649-2
17. Pang J, Salim A, Lee VJ, Hibberd ML, Chia KS, Leo YS, Lye DC: Diabetes with hypertension as risk factors for adult dengue hemorrhagic fever in a predominantly dengue serotype 2 epidemic: a case control study. *PLoS Negl Trop Dis* 2012; 6(5):e1641. doi: 10.1371/journal.pntd.0001641