

CASE REPORT

Off-label Use of Itolizumab in Patients with COVID-19 ARDS: Our Clinical Experience in a Dedicated COVID Center

Poonam Kumari¹, Amarjeet Kumar², Chandni Sinha³, Ajeet Kumar⁴, Prabhat K Singh⁵, SK Arun⁶

ABSTRACT

Severe acute respiratory syndrome coronavirus 2 has affected millions of people worldwide. This pandemic requires newer medical management strategies to control the morbidity and mortality associated with the disease. Several approaches, including global targeting of inflammation or neutralizing a single key inflammatory mediator, are being employed to cope with cytokine storms in COVID-19. The role of anti-inflammatory biologics, such as acalabrutinib, tocilizumab, anakinra, and itolizumab can become relevant. Itolizumab is a humanized recombinant immunoglobulin G1 monoclonal antibody. It targets the extracellular, scavenger receptor cysteine-rich (SRCR) distal domain 1 of CD6 and is responsible for priming, activation, and differentiation of T-cells. Itolizumab has been approved by the Drug Controller General of India for the treatment of COVID 19 in India. Here, we shared our clinical experience of 20 patients having moderate acute respiratory distress syndrome (ARDS) due to COVID-19 on treatment with itolizumab. We observed the mortality benefit with single-dose itolizumab (1.6 mg/kg) in patients having moderate COVID-19 ARDS.

Keywords: Acute respiratory distress syndrome (ARDS), COVID-19, Cytokine storm, Itolizumab, Monoclonal antibody.

Indian Journal of Critical Care Medicine (2021): 10.5005/jp-journals-10071-23787

HIGHLIGHTS

- Fifteen percent of COVID-19 patients who develop acute respiratory distress syndrome within a week after the appearance of symptoms.
- Cytokine storm is a life-threatening acute systemic inflammatory syndrome, which is characterized by multiorgan failure and fever.
- Itolizumab is an anti-CD6 receptor monoclonal antibody which seems to be the possible strategy for patients with cytokine storm.
- We used Itolizumab off-label in patients with severe COVID-19 from August to September 2020 based on its hypothetical benefit and report the experience in our institute AIIMS, Patna.

INTRODUCTION

Severe acute respiratory syndrome (SARS) coronavirus 2 (CoV-2) has affected millions of people worldwide. This pandemic requires newer medical management strategies to control the morbidity and mortality associated with the disease. About 15 to 20% of the affected patients develop pulmonary symptoms, such as breathing difficulty, and require hospital admission for oxygen therapy and supportive care.¹ There is a marked increase in the level of cytokines (e.g., interleukin(IL)-1B, IL-6, and IL-12) and chemokines and this is referred to as "cytokine storm."² There is an elevation of other markers of inflammation, coagulation, and organ damage, such as C-reactive protein (CRP), D-dimer, lactate dehydrogenase, ferritin, and troponin-I. Hence, anti-inflammatory agents, such as acalabrutinib, tocilizumab, anakinra, and itolizumab might have a role to play in this disease.³⁻⁸ Itolizumab is a humanized recombinant immunoglobulin G1 monoclonal antibody. It targets extracellular, scavenger receptor cysteine-rich distal domain 1 of

^{1,3,4,6}Department of Anaesthesiology, AIIMS Patna, Patna, Bihar, India

²Department of Trauma and Emergency, AIIMS Patna, Patna, Bihar, India

⁵Director, AIIMS Patna, Patna, Bihar, India

Corresponding Author: Amarjeet Kumar, Department of Trauma and Emergency, AIIMS Patna, Patna, Bihar, India, Phone: +91 09570890646, e-mail: amarjeetdmch@gmail.com

How to cite this article: Kumari P, Kumar A, Sinha C, Kumar A, Singh PK, Arun SK. Off-label Use of Itolizumab in Patients with COVID-19 ARDS: Our Clinical Experience in a Dedicated COVID Center. *Indian J Crit Care Med* 2021;25(4):467–469.

Source of support: Nil

Conflict of interest: None

CD6 and is responsible for priming, activation, and differentiation of T-cells.⁹ Drug Controller General of India (DCGI) has approved itolizumab for the treatment of COVID-19 in India. Here, we share our single-center clinical experience of off-label use of this drug in 20 patients with acute respiratory distress syndrome (ARDS) due to COVID-19.

METHODS

This is our single-center clinical experience of patients infected with COVID-19 and treated with itolizumab between August and September 2020 at All India Institute of Medical Sciences, Patna. We recorded various clinical and biochemical parameters every day from the day of admission to 14 days after administration of itolizumab. Clinical parameters, such as respiratory rate and oxygen supplementation, were measured daily. Laboratory parameters [CRP, lymphocytes, aspartate aminotransferase (AST), and alanine

aminotransferase (ALT)] were measured at the clinician's discretion and collected from baseline to day 14.

Due to high costs, lack of knowledge, and safety risks, a multidisciplinary ethics committee validated the off-label use of itolizumab in moderately severe patients. Patient consent was requested and marked in the patients' records.

The patients who received itolizumab had the following criteria:

- Age between 18 and 60 years
- Virological diagnosis of SARS-CoV2 infection (polymerase chain reaction)
- Patients who were suffering from moderate COVID-19 ARDS (as defined by PaO₂/FiO₂ ratio of <200, or more than 25% deterioration from the immediate previous value within 24 hours in patients having more than PaO₂/FiO₂ >200).
- Serum ferritin level ≥ 400 ng/mL at baseline or IL-6 levels greater than four times the upper limit of normal.

Exclusion criteria for injection itolizumab included: patients with a history of severe allergic reactions to monoclonal antibodies, history of active or latent tuberculosis (TB) infection, patients on oral immunosuppressive drugs within the past 6 months, pregnant or breastfeeding patients, or those with a positive pregnancy test in a pre-dose examination. Patients with a known history of hepatitis B, hepatitis C, or HIV, absolute neutrophil count <1000/mm³, platelets <50000/mm³, absolute lymphocyte count <500/mm³, very severe ARDS PaO₂/FiO₂ ≤100 with positive end-expiratory pressure (PEEP) ≥5, and ARDS due to other secondary infections/bacterial sepsis (high procalcitonin value) were also not administered the drug.

Patients of itolizumab group received the drug at a dose of 1.6 mg/kg. Itolizumab was dissolved in 250 mL of 0.9% normal saline. The infusion was started at a rate of 25 mL in the first hour and given over a time period of 4 to 6 hours. During infusion, vitals of the patient were monitored. Premedication with hydrocortisone 100 mg and pheniramine 30 mg intravenous was given 30 minutes before infusion. The second dose of the drug was repeated according to clinical evaluation, response to the first dose, and lab parameters. All patients admitted to the intensive care unit (ICU) received standard baseline treatments as per SOP AIIMS, Patna. Antibiotics were given as per antibiogram report. During hospitalization, these patients required oxygen therapy in the form of nonrebreathing mask (NRM), high-flow nasal cannula (HFNC), continuous positive airway pressure (CPAP), or invasive (i.e., mechanical ventilation).

In this clinical experience, we used itolizumab in a very small subset of patients admitted in COVID ICU. This allowed us to compare the effect of itolizumab versus control. We compared the parameters in both groups of these patients: one treated with itolizumab and the other with a relatively similar clinical impairment who had not been treated with any other immunomodulators.

We have limited crude raw data of patients who have not received itolizumab. Bias could not be eliminated in the groups due to presence of sicker patients with comorbidities in the control group.

RESULTS

Patients' Outcome

From August 1, 2020, to September 30, 2020, totally 150 patients were included in our study. Of these 150, 20 received injection

itolizumab (1.6 mg/kg) following the fulfillment of inclusion criteria. Among these patients who received itolizumab, seven patients were not survived and 1-month mortality rate was 35%. Among these patients who have not received itolizumab were 130 and 1-month mortality was 60%.

Adverse Effects

We detected infusion-related adverse reactions in eight patients; six patients have mild shivering and other two patients noted tachycardia and oxygen desaturation, for which they required increment of oxygen support.

DISCUSSION

There are various drugs like tocilizumab, itolizumab, and anakinra used in the treatment of COVID-19. Among these, itolizumab is a new biological agent that has been approved in India for the treatment of psoriasis.¹⁰ It acts by immunomodulating T-effector function and its migration to the inflammation site, sparing Tregs, and preserving the antiviral response. It acts upstream at the Th1 and Th17 pathways,^{9,11} i.e., at the T-effector cells. Thus, it decreases the release of multiple cytokines and cell signaling transduction factors primarily involving the Th17 and Th1 pathways. Other drugs, such as tocilizumab or anakinra, block cytokines at a downstream level. Itolizumab also binds to CD6 receptor and blocks activated leukocyte cell adhesion molecule-mediated T-cell activation. For this reason, it is used in the treatment of COVID-19 patient clinical trial done in Cuba.¹² Tocilizumab was also used in the treatment of COVID-19 patients with moderate-to-severe ARDS.¹³ Albertini et al.¹⁴ in an observational study over off-label use of tocilizumab concluded that tocilizumab appears to be effective in decreasing oxygen withdrawal time and respiratory failure symptoms, avoiding mechanical ventilation, and stopping the biological inflammatory process of COVID-19 pneumonia in hospitalized patients, especially in those with >50% of initial lung lesions on computed tomography scan. It is important to mention that IL-6 is not the appropriate marker of disease improvement as the soluble IL-6 receptor immune complex has a long half-life. It remains in the serum for a long time because of its long half-life after tocilizumab administration. IL-6 levels might also increase after tocilizumab administration because of less consumption of IL-6.¹³

We observed mortality benefit (35 vs 60%) in moderate COVID-19 ARDS patients who have received single-dose itolizumab 1.6 mg/kg in comparison to patients who have not received immunomodulator therapy (tocilizumab/itolizumab) in the course of COVID-19 disease. In India, the mortality rate in moderate-to-severe COVID-19 pneumonia is 42%.¹⁵ Tocilizumab was another immunomodulator used at our center. In our experience, despite giving two doses within 12 hours we have found no improvement. We also observed cost benefit in patients receiving itolizumab, as a single dose was sufficient.

A randomized, controlled, open-label study was conducted by Biocon at four hospitals in India, in moderate-to-severe COVID-19 ARDS patients. As reported by Biocon, the mortality benefit observed in the itolizumab arm was statistically significant than the only supportive care group.

Limitations

Here, we have shared our clinical experience with moderate COVID ARDS patients on treatment with itolizumab. We have not compared the clinical outcome of patients on treatment with

other drugs like tocilizumab. It would be difficult to comment on or to speculate that itolizumab has better survival benefits without a well-conducted randomized controlled trial, comparing various modalities. Nonetheless, this clinical experience does make us consider itolizumab the treatment option. We have not included patients with mild and severe ARDS ($\text{PaO}_2/\text{FiO}_2 \leq 100$ with $\text{PEEP} \geq 5$). Due to unequal distribution of comorbidities among patients, we acknowledge that our results should be interpreted with caution as significant heterogeneity exists within these factors.

CONCLUSION

We observed mortality benefit with single-dose itolizumab (1.6 mg/kg) in patients having moderate COVID-19 ARDS. A well-structured comparative study is required to validate our result and to find superiority over tocilizumab. Use and recommendations for itolizumab in COVID-19 moderate/severe disease should be taken with caution as not only the supportive data are lacking but also long-term outcomes of itolizumab in COVID-19 diseases are yet to be known.

ORCID

Poonam Kumari  <https://orcid.org/0000-0003-0118-2187>

Amarjeet Kumar  <https://orcid.org/0000-0002-4272-5750>

Chandni Sinha  <https://orcid.org/0000-0002-4107-2671>

Ajeet Kumar  <https://orcid.org/0000-0002-1464-6684>

Prabhat K Singh  <https://orcid.org/0000-0003-1888-4679>

Arun SK  <https://orcid.org/0000-0002-9788-3350>

REFERENCES

- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061–1069. DOI: 10.1001/jama.2020.1585.
- Coperchini F, Chiovato L, Croce L, Magri F, Rotondi M. The cytokine storm in COVID-19: An overview of the involvement of the chemokine/chemokine-receptor system. *Cytokine Growth Factor Rev* 2020;53:25–32. DOI: 10.1016/j.cytogfr.2020.05.003.
- Roschewski M, Lionakis MS, Sharman JP, Roswarski J, Goy A, Monticelli MA, et al. Inhibition of Bruton tyrosine kinase in patients with severe COVID-19. *Sci Immunol* 2020;5(48):eabd0110. DOI: 10.1126/sciimmunol.abd0110.
- Alijotas-Reig J, Esteve-Valverde E, Belizna C, Selva-O'Callaghan A, Pardos-Gea J, Quintana A, et al. Immunomodulatory therapy for the management of severe COVID-19. Beyond the anti-viral therapy: a comprehensive review. *Autoimmun Rev* 2020;19(7):102569. DOI: 10.1016/j.autrev.2020.102569.
- Marovich M, Mascola JR, Cohen MS. Monoclonal antibodies for prevention and treatment of COVID-19. *JAMA* 2020;324(2):131–132. DOI: 10.1001/jama.2020.10245.
- Bughani U, Saha A, Kuriakose A, Nair R, Sadashivarao RB, Venkataraman R, et al. T cell activation and differentiation is modulated by a CD6 domain 1 antibody Itolizumab. *PLoS One* 2017;12(7):e0180088. DOI: 10.1371/journal.pone.0180088.
- Guaraldi G, Meschiari M, Cozzi-Lepri A, Milic J, Tonelli R, Menozzi M, et al. Tocilizumab in patients with severe COVID-19: a retrospective cohort study. *Lancet Rheumatol* 2020;2(8):e474–e484. DOI: 10.1016/S2665-9913(20)30173-9.
- Cavalli G, De Luca G, Campochiaro C, Della-Torre E, Ripa M, Canetti D, et al. Interleukin-1 blockade with high-dose anakinra in patients with COVID-19, acute respiratory distress syndrome, and hyperinflammation: a retrospective cohort study. *Lancet Rheumatol* 2020;2(6):e325–e331. DOI: 10.1016/S2665-9913(20)30127-2.
- Nair P, Melarkode R, Rajkumar D, Montero E. CD6 synergistic co-stimulation promoting proinflammatory response is modulated without interfering with the activated leucocyte cell adhesion molecule interaction. *Clin Exp Immunol* 2010;162(1):116–130. DOI: 10.1111/j.1365-2249.2010.04235.x.
- Gupta A, Sharma YK, Deo K, Kothari P. Severe recalcitrant psoriasis treated with itolizumab, a novel anti-CD6 monoclonal antibody. *Indian J Dermatol Venereol Leprol* 2016;82(4):459–461. DOI: 10.4103/0378-6323.181466.
- Anand A, Assudani D, Nair P, Krishnamurthy S, Deodhar S, Arumugam M, et al. Safety, efficacy and pharmacokinetics of T1h, a humanized anti-CD6 monoclonal antibody, in moderate to severe chronic plaque psoriasis – results from a randomized phase II trial. (96.13). *J Immunol* 2010 Apr 1;184(1 Supplement) 96.13 [Phase 2 trial of Itolizumab in moderate to severe chronic plaque psoriasis].
- Treatment of patients with severe SARS-CoV-2 pneumonia with the anti-CD6 monoclonal antibody itolizumab-full text-WHO-ICTRP [Internet] [Cited 2020 Jul 16]. Available from: <https://apps.who.int/trialsearch/Trial2.aspx?TrialID=RPCEC00000311>.
- Xu X, Han M, Li T, Sun W, Wang D, Fu B, et al. Effective treatment of severe COVID-19 patients with tocilizumab. *Proc Natl Acad Sci U S A* 2020;117(20):10970–10975. DOI: 10.1073/pnas.2005615117.
- Albertini L, Soletchnik M, Razurel A, Cohen J, Bidegain F, Fauvelle F, et al. Observational study on off-label use of tocilizumab in patients with severe COVID-19. *Eur J Hosp Pharm* 2020;0:1–7. DOI: 10.1136/ejhpharm-2020-002414.
- Fewer COVID-19 deaths in ICU suggest hospital care improving: study; 2020 [Accessed: September 20, 2020]. Available from: <https://www.ndtv.com/world-news/coronavirus-fewer-covid-19-deaths-in-icu-suggest-hospital-care-improving-study-2264244>.