

Effect of Percutaneous Tracheostomy on Optic Nerve Sheath Diameter (TONS Trial)

Ilenia Di Paola¹, Mario Graziano², Palmiro Cornetta³

Keywords: Anesthetists, A-scan, B-scan, Optic nerve sheath diameter, Ophthalmology, Surgery.
Indian Journal of Critical Care Medicine (2022): 10.5005/jp-journals-10071-24208

To the Editor

We read with great interest the article by Kapoor et al. concerning the effect of percutaneous tracheostomy (PCT) on optic nerve sheath diameter (ONSD).¹

We congratulate the authors for their very interesting study, but we would like to make some comments.

In this paper, the authors found an increase in the ONSD during elective PCT in neurocritical ill patients.

Geng et al. described an increase in the ONSD in patients that underwent anesthesia with propofol compared to patients that underwent anesthesia with sevoflurane.²

As the authors utilized propofol and sevoflurane for the anesthesia, we were wondering if the increase in ONSD is really surgical procedure related or it could be related to the anesthesia.

Maybe it could be useful for further studies to check if utilizing different anesthetic procedures, same results can be obtained.

Another point is that the cutoff value observed in this study was considerably different from previously reported studies. The reason for this could be related to the use of B-scan that is affected by several limitations due to the lack of standardization and presence of artifacts.^{3,4} Unfortunately, even taking into account the advice to image the central retinal artery with color Doppler these problems cannot be overcome.

For this reason, to prevent these artifacts, we suggest, in case of future studies, to use the so-called standardized A-scan technique.⁵

ORCID

Ilenia Di Paola <https://orcid.org/0000-0003-1444-6663>

Mario Graziano <https://orcid.org/0000-0002-5477-5042>

Palmiro Cornetta <https://orcid.org/0000-0001-9831-1955>

^{1,2}Department of Medicine, Surgery and Dentistry, "Scuola Medica Salernitana", University of Salerno, Baronissi, Salerno, Italy

³ASL Salerno, Salerno, Italy

Corresponding Author: Mario Graziano, Department of Medicine, Surgery and Dentistry, "Scuola Medica Salernitana", University of Salerno, Baronissi, Salerno, Italy, e-mail: mgraziano@unisa.it

How to cite this article: Di Paola I, Graziano M, Cornetta P. Effect of Percutaneous Tracheostomy on Optic Nerve Sheath Diameter (TONS Trial). *Indian J Crit Care Med* 2022;26(5):651.

Source of support: Nil

Conflict of interest: None

REFERENCES

1. Kapoor I, Wanchoo J, Mahajan C, Singhal V, Roy H, Kumar S, et al. Effect of percutaneous tracheostomy on optic nerve sheath diameter [TONS Trial]. *Indian J Crit Care Med* 2021;25(4):382–387. DOI: 10.5005/jp-journals-10071-23783.
2. Geng W, Chen C, Sun X, Huang S. Effects of sevoflurane and propofol on the optic nerve sheath diameter in patients undergoing laparoscopic gynecological surgery: a randomized controlled clinical studies. *BMC Anesthesiol* 2021;21(1):30. DOI: 10.1186/s12871-021-01243-7.
3. De Bernardo M, Vitiello L, Rosa N. Ultrasound optic nerve sheath diameter evaluation in patients undergoing robot-assisted laparoscopic pelvic surgery. *J Robot Surg* 2019;13(5):709–710. DOI: 10.1007/s11701-019-00966-7.
4. De Bernardo M, Vitiello L, Rosa N. Optic nerve evaluation in idiopathic intracranial hypertension. *Am J Neuroradiol* 2019;40(7):E36. DOI: 10.3174/ajnr.A6091.
5. Rosa D, Graziano M, Di Paola I. Evaluation of intracranial pressure during neural laser dissection. *Pain Physician J* 2022;25(2):E414. PMID: 35323006.