

Optic Nerve Sheath Diameter in Hyponatremia: A Closer Look

Jes Jose¹, Varun Suresh², Rohan Magoon³

Received on: 14 April 2023; Accepted on: 14 April 2023; Published on: 31 May 2023

Keywords: Hyponatremia, Optic nerve sheath diameter, Ultrasonography.

Indian Journal of Critical Care Medicine (2023): 10.5005/jp-journals-10071-24454

Dear Editor,

The research study by Uttanganakam et al.¹ brings to light a significant decrease in optic nerve sheath diameter (ONSD) after treatment of hyponatremia compared to pretreatment levels (mean right eye ONSD = 5.81 ± 0.58 mm [95% confidence interval (CI): 0.27–0.59, $p < 0.001$], and left eye ONSD = 5.79 ± 0.56 mm [95% CI: 0.29–0.64, $p < 0.001$] after hyponatremia correction; compared to mean right eye ONSD = 6.24 ± 0.71 mm and left eye ONSD = 6.26 ± 0.64 mm at presentation to the emergency department). However, the ONSD measurements failed to predict moderate to severe hyponatremia on receiver operating characteristic curve (ROC) analysis in the index prospective cohort. The author's attempt to extend the point of care utility of ultrasonographic measured ONSD to the emergency department deserves applause, though herewith we appraise a few facets to consider while extrapolating sub-millimetric variations in ONSD to clinical outcome.²

Uttanganakam et al. measured ONSD in the transverse plane with a high-frequency linear ultrasound probe using B-mode with the patient eyes closed, using hypoechoic lines behind the eye globe as reference. In B-scan, the ultrasonographic waves reach the optic nerve in a non-perpendicular array and hence measurements are predisposed to "scattering" and "blooming effect" artifact.³ Despite the use of an ophthalmic setting Uttanganakam et al. contemplated, ONS wall brightness varies with 'gain' adjustments in B-scan. That said the index measurement being ONSD, variations in microns can affect the accuracy and precision of readings.⁴ Therefore, we suggest that A-scan ultrasound (wherein the ONS is easily noticeable as hyperreflective spikes), be used especially so in the emergency department cohort of patients as time constraints and quicker decision-making are pertinent here.

That, 44.4% of cases in the index cohort had vomiting leaves the readers intrigued by the extent to which dehydration could have influenced ONSD measurements in hypovolemic hyponatremia. It is appreciated that the authors have carefully excluded hypervolemic hyponatremia, yet hypovolemia can alter cerebrospinal fluid production and dynamics consequently affecting ONSD measurements. Though Uttanganakam et al. 'practiced' ONSD imaging in 80 individual measurements prior to the study to avoid interobserver variability in measurements, how the authors 'blinded' their investigators to the nature of the image procured (before and after sodium correction) has been left unaccounted. Further, we appreciate that Uttanganakam et al. has carefully excluded head injury patients from their research wherein septations and ONS compartmentalization can mandate separate correlation of right and left ONSD measurements to the studied outcome, nevertheless, we believe

¹Department of Cardiac Anesthesiology, Sri Jayadeva Institute of Cardiovascular Sciences and Research, Bengaluru, Karnataka, India

²Department of Anesthesia and Intensive Care, Jaber Al Ahmad Al Sabah Hospital, Kuwait-Arabian Gulf

³Department of Anaesthesia, Atal Bihari Vajpayee Institute of Medical Sciences (ABVIMS) and Dr. Ram Manohar Lohia Hospital, New Delhi, India

Corresponding Author: Varun Suresh, Department of Anesthesia and Intensive Care, Jaber Al Ahmad Al Sabah Hospital, Kuwait-Arabian Gulf, Phone: +9041426743, e-mail: varunsureshpgi@gmail.com

How to cite this article: Jose J, Suresh V, Magoon R. Optic Nerve Sheath Diameter in Hyponatremia: A Closer Look. *Indian J Crit Care Med* 2023;27(6):452.

Source of support: Nil

Conflict of interest: None

that the average of either side ONSD can be better representative in the correlation and ROC analysis of the Uttanganakam et al. hyponatremia cohort of cases.⁵

ORCID

Jes Jose  <https://orcid.org/0000-0003-1734-9519>

Varun Suresh  <https://orcid.org/0000-0003-2521-1149>

Rohan Magoon  <https://orcid.org/0000-0003-4633-8851>

REFERENCES

1. Uttanganakam S, Hansda U, Sahoo S, Shaji IM, Guru S, Topno N, et al. Sonographic optic nerve sheath diameter as a guide for correction of hyponatremia in the emergency department: A cross-sectional study. *Indian J Crit Care Med* 2023;27(4):265–269. DOI: 10.5005/jp-journals-10071-24438.
2. Arunachala S. Optic nerve sheath diameter and sodium levels: A friend of a friend is still a stranger. *Indian J Crit Care Med* 2023;27(4):235–236. DOI: 10.5005/jp-journals-10071-24444.
3. Cornetta P, Marotta G, De Bernardo M, Vitiello L, Rosa N. Ultrasound and optic neuritis. *Am J Emerg Med* 2019;37(8):1598. DOI:10.1016/j.ajem.2019.02.001.
4. Stevens RRF, Gommer ED, Aries MJH, Ertl M, Mess WH, Huberts W, et al. Optic nerve sheath diameter assessment by neurosonology: A review of methodologic discrepancies. *J Neuroimaging* 2021;31(5):814–825. DOI:10.1111/jon.12906.
5. Suresh V, Sharma S, Aggarwal A. Correlation between ultrasonographic optic nerve sheath diameter and intracranial pressure in patients with aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2020;33(3):860–861. DOI: 10.1007/s12028-020-01087-y.